



Review Article

Asparagus racemosus (Shatavari): A Review on *Asparagus racemosus* (Shatavari): From Traditional Medicine to Modern Therapeutics

Abstract

Asparagus racemosus (Shatavari) is one of the most valuable medicinal plants in Traditional Medicine. ~~Ayurveda~~ and has been widely used for centuries to manage reproductive, gastrointestinal, neurological, and inflammatory disorders. Growing scientific interest has prompted extensive investigations into its phytochemical composition and pharmacological activities. This review provides a comprehensive overview of the botanical characteristics, phytochemical constituents, traditional applications, and therapeutic potential of *A. racemosus*. The plant is rich in steroidal saponins, particularly shatavarins, along with flavonoids, alkaloids, polyphenols, essential fatty acids, vitamins, and trace minerals that collectively contribute to its diverse biological activities. Experimental and clinical studies demonstrate that *A. racemosus* exhibits antioxidants, anti-inflammatory, immunomodulatory, adaptogenic, antimicrobial, antiulcer, antidiabetic, hepatoprotective, neuroprotective, galactagogue, aphrodisiac, and anticancer properties. In addition, emerging evidence supports its beneficial effects on female reproductive health, cognitive function, stress reduction, and metabolic regulation. Despite substantial preclinical evidence, further well-designed clinical trials, together with standardized phytochemical profiling, pharmacokinetic investigations, and long-term safety evaluations, are needed to establish therapeutic efficacy and safety and support its integration into evidence-based medicine. ~~Future research incorporating advanced analytical techniques, molecular docking, systems biology, and nanotechnology based drug delivery may accelerate the development of standardized~~

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Commented [Editor1]: Better replace "Ayurveda" with "in traditional medicine"

Formatted: Font color: Blue

Ali et al.

(e-ISSN: 3078-6975)

phytopharmaceutical products derived from *A. racemosus*. Overall, this review highlights the significant medicinal value of *A. racemosus* and underscores its potential as a promising natural therapeutic agent for the prevention and management of various human diseases.

Keywords: *Asparagus racemosus*; Shatavari; Medicinal plant; Phytochemicals; Steroidal saponins; Shatavarins; Pharmacological activities; Antioxidant; Anti-inflammatory; Immunomodulatory; Adaptogen; Ayurveda.

Commented [Editor2]: You should mention comprehensive in vivo studies and clinical studies for therapeutic efficacy and safety studies are required.

Formatted: Font color: Blue

Introduction

Medicinal plants have a long history of use in treating a wide range of human ailments (Saggar et al., 2022). Although the rise of modern allopathic medicine led to a decline in traditional medicinal practices, recent decades have witnessed a global resurgence. This renewed interest is driven by several factors, including the side effects and toxicity associated with many synthetic drugs, the emergence of multidrug-resistant microorganisms, and the inability of modern medicine to provide effective cures for numerous chronic diseases. Today, more than 70% of the population in developing countries relies on traditional medicinal systems, often referred to as complementary or alternative medicine for primary healthcare (Ekor, 2014).

Plants used by indigenous communities continue to serve as valuable sources of novel pharmaceuticals (Gurib-Fakim, 2006). For example, traditional Native American medicine has contributed unique approaches to cardiovascular disease management that complement modern treatments (Fabricant and Farnsworth, 2001). Several widely used drugs, such as the anticancer agent’s vinblastine and taxol, and the antimalarial compound artemisinin, are derived from medicinal plants. A South African herbal remedy traditionally used for respiratory infections has also shown promise in treating bronchitis. Recent studies further highlight the pharmacological potential of medicinal plants, including their anticancer, antioxidant, and antibacterial activities (Atanasov et al., 2015; Faruquee et al., 2026). These examples underscore the growing global interest in medicinal plants as accessible, affordable, and culturally relevant healthcare resources.

Bangladesh has a rich heritage of traditional medicinal practices. Traditional healers continue to serve as primary healthcare providers for many rural communities where access to modern healthcare remains limited. Recent evidence also indicates that self-medication and reliance on non-professional healthcare advice remain common in Bangladesh, emphasizing the continued importance of medicinal plants and traditional healthcare practices in primary health management (Kuate et al., 2008; Shahjalal et al., 2025). Because the selection and use of medicinal plants vary significantly across regions, systematic surveys are essential to document the diversity of medicinal plant use throughout the country. The present study was conducted in the Chalna area of Dacope Upazila in Khulna District, located near the Sundarbans forest. In this region, modern healthcare facilities are scarce, and residents depend heavily on traditional practitioners for medical care.

Asparagus racemosus (commonly known as satavar, shatavari, or shatamull) is one such medicinal plant widely used in traditional systems (Alok et al., 2013). Native to India, Nepal, Sri Lanka, and the Himalayan region, the plant grows one to two meters tall (Singh and Geetanjali, 2016). In Ayurveda, it is revered as the “Queen of Herbs” for its rejuvenating properties and its association with love, vitality, and devotion. Shatavari is considered the primary female rejuvenative tonic in Ayurveda, much like *Withania somnifera* is for males. The name “Shatavari,” meaning “she who possesses a hundred husbands,” reflects its traditional use in supporting female reproductive health.

In Ayurveda, *A. racemosus* is described as an effective remedy for chronic fever and internal heat. Its strong association with female reproductive health is well documented in classical Ayurvedic texts (Kushwah et al., 2018). Modern Ayurvedic practice attributes numerous therapeutic properties to the roots of *A. racemosus*, including antispasmodic, stomachic, aphrodisiac, galactagogue, astringent, antidiarrheal, antidyenteric, laxative, anticancer, anti-inflammatory, blood-purifying, antitubercular, and antiepileptic effects. Recent pharmacological studies have expanded this understanding, demonstrating

anticancer, antioxidant, and antibacterial activities, potential therapeutic effects in epilepsy and memory dysfunction, and promising neuroprotective activity relevant to Alzheimer’s disease (Shaji et al., 2025). Additional research highlights its food-based applications and functional properties, further increasing its global demand (Akhtar et al., 2024). First described botanically in 1799, the plant has seen increasing demand due to its wide range of medicinal applications. However, destructive harvesting practices, habitat loss, and deforestation have led to its classification as an endangered species in the wild.

Formatted: Font color: Blue

Botanical Identification and Description of the plant

The genus *Asparagus* has been recently moved from the subfamily *Asparagae* in the family *Liliaceae* to a newly created family *Asparagaceae*. The *Asparagus* genus is of medicinal importance because of the presence of steroidal saponins and sapogenins in various parts of the plant (Shao et al., 1997; Dawid and Hofmann, 2012). *Asparagus* is the Greek word for “stalk” or “shoot.” About 300 species of *Asparagus* are known to occur in the world. Some of the European species to be mentioned are *A. officinalis*, *A. sprengeri*, and *A. acutifolius*. Among the several species of *Asparagus*, *Asparagus racemosus*, *Asparagus gonacledes*, and *Asparagus adsendens* grow in India and are most commonly used in indigenous medicine (Singh and Geetanjali, 2016). *Asparagus racemosus* is the one most commonly used in traditional medicine due to its wide pharmacological potential (Banerjee et al., 2025).

Kingdom:	Plantae
Subkingdom:	Tracheobionta
Division:	Magnoliophyta
Class:	Magnoliopsida
Order:	Asparagales
Family:	Asparagaceae
Genus:	Asparagus
Species:	racemosus

Botanical name: *Asparagus racemosus*

Asparagus racemosus is commonly called Satavari, Satawar or Satmuli in Hindi, Satavari in Sanskrit, Shatamuli in Bengali, Shatmuli in Marathi, Satawari in Gujarati, Toala-gaddalu or Pilli-gaddalu in Telugu, Shimaishadavari or Thanner Vittan Kizhangu or Inli-chedi in Tamil, Chatavali in Malayalam, Majjigegadde or Aheruballi in Kannada, Kairuwa in Kumaon, Narbodh or Satmooli in Madhya Pradesh, and Norkanto or Satawar in Rajasthan (Thakur et al., 2015).

Formatted: Font color: Blue

The plant grows throughout the tropical and subtropical parts of India up to an altitude of 1500 m.


Roots: The plant is a spinous under-shrub, with tuberous, short rootstock bearing numerous succulent tuberous roots (30–100 cm long and 1–2 cm thick) that are silvery white or ash-colored externally and white internally. These roots are the part that finds use in various medicinal preparations (Ms et al., 2018; Singh et al., 2023). **Stem:** The stem is woody, climbing, whitish grey or brown colored with small spines. **Flowers:** The plant flowers during February–March, leaving a mild fragrance in its surroundings. **Fruits:** By the end of April, fruits are visible, with attractive red berries.

A thorny, climbing shrub with woody stems. Leaves are reduced to minute scales and spines. Fruits are globular. Roots are succulent and tuberous which are tapering at both ends. Subshrubs hermaphroditic, stems climbing, branched to 2 m, branches usually distinctly striate-ridged, ridges ± cartilaginous denticulate. Cladodes in fascicles of 3–6(–8), linear, 1-2.5 cm × 1 mm, flat, midvein distinct. Leaf spurs spinescent; spine straight or subrecurved, 1.5–2 cm on main stems, 5–10 mm on branches, woody, sharp. Inflorescences developing after cladodes, axillary, each a many-flowered raceme or panicle 1–4 cm, bracts ~1 mm. Pedicel 1.5–3 mm, slender articulate at middle (Frawley, 1989; Thakur et al., 2015; Ms et al., 2018; Gogte, 2022)

Phytochemical constituents Chemical Constituents

Formatted: Font color: Blue


The major bioactive constituents of *Asparagus* are a group of steroidal saponins. This plant also contains vitamins A, B1, B2, C, E, Mg, P, Ca, Fe, and folic acid. Other primary chemical constituents of *Asparagus* are essential oils, asparagine, arginine, tyrosine, flavonoids (kaempferol, rutin), resin, and tannin [48]. Shatavarin IV is a glycoside of sarsasapogenin having two molecules of *Asparagus* rhamnose and one molecule of glucose. The major bioactives (chemical constituents) of *Asparagus* species are shown in sarsasapogenin and shatavarin I–IV, which are present in roots, leaves, and fruits of *Asparagus* species. Synthesis of sarsasapogenin in the callus culture of *A. racemosus* was also reported (Bopana and Saxena, 2007). A new isoflavone, 8-methoxy-5,6,4'-trihydroxyisoflavone-7-O-β-D-glucopyranoside, was also reported from *A. racemosus* previously (Saxena and Chourasia, 2001). The isolation and characterization of a polycyclic alkaloid called asparagamine (Sekine et al., 1994), a new 9,10-dihydrophenanthrene derivative named racemosol, and kaempferol were also isolated from the ethanolic root extract of *A. racemosus* (Sekine et al., 1997). Oligofurostanosides (curillins G and H) and spirostanosides (curilloside G and H) have been isolated from the roots, and sarsasapogenin from leaves of *A. curillus*. Recent research has expanded the phytochemical profile of *A. racemosus*, confirming additional saponins, flavonoids, and phenolic antioxidants that support its traditional medicinal applications (Saxena, 2025). Modern reviews have emphasized its adaptogenic, immunomodulatory, and reproductive-supportive effects, linking these activities to its diverse saponin and isoflavonoid content (Mateen et al., 2025). Ethnopharmacological studies published between 2016 and 2025 have further validated its classical Ayurvedic uses, particularly in female reproductive health and gastrointestinal protection (Ahmed and Urooj, 2010). Clinically oriented investigations also suggest that *A. racemosus* may enhance fertility due to its phytoestrogenic constituents (Oyovwi et al., 2025). Advanced phytochemical profiling using in silico ADMET and antioxidant modeling has identified new bioactive molecules in the roots, reinforcing its therapeutic potential (Chikhale et al., 2021a).



R=H, Sarsasapogenin

R=Glu[(4-1)Rha](2-1).Glu, Shatavarin

Fig 15: Structures of sarsasapogenin and its glycosides.



R=Glu(6-1), Adscendin A

R=Glu[(4-1)Rha](6-1)Rha, Adscendin B

R=Glu(2-1)Glu, Asparanin A

R=Glu[(2-1)Glu](4-1)Rha, Asparanin B

R=Glu(4-1)Ara(6-1)Rha, Asparanin C

R=Glu[(2-1)Rha](4-1)Ara, Curillin G

R=Glu[(2-1)Glu](4-1)Ara, Curillin H

Fig 26: Isolated compounds from *Asparagus* species.

The structural complexity of saponins results in a number of physical, chemical, and biological properties. Saponins are usually amorphous substances with high molecular weights. These are soluble in water and produce foam, but organic solvents such as chloroform, acetone, and ether inhibit their foaming property. Solubility of saponins is also influenced by the properties of the solvent (temperature, composition, and pH), whereas water, alcohols (methanol, ethanol), and aqueous alcohols are the most common extraction solvents for saponins. Due to the presence of a lipid-soluble aglycone and water-soluble sugar chain in their structure (amphiphilic nature), saponins are surface-active compounds with detergent, wetting, emulsifying, and foaming properties. In aqueous solutions, surfactants form micelles above a critical concentration called the critical micelle concentration.

Saponins possess a variety of biological properties, namely antioxidant, immunostimulant, antihepatotoxic, antibacterial, and useful in diabetic retinopathy, anticarcinogenic, antidiarrheal,

antiulcerogenic, antioxytotic, and reproductive activities. Many saponins are known to be antimicrobial, to inhibit mould, and to protect plants from insects. They may be considered as a defense system and have been included in a large group of protective molecules found in plants named phytoanticipins or phytoprotectants. Saponin-rich plants have been found to improve growth, feed efficiency, and health in ruminants (Zhang *et al.*, 2019).

Phytochemical constituents

Steroidal saponins, known as shatavarins. Shatavarin I to VI are present. Shatavarin I is the major glycoside with 3-glucose and rhamnose moieties attached to sarsapogenin (Hayes *et al.*, 2006; Guo *et al.*, 2023). Oligospirostanoside referred to as Immunoside (Lomelino *et al.*, 2017). Polycyclic alkaloid Asparagamine A, a cage-type pyrrolizidine alkaloid (Sekine *et al.*, 1995; Potduang *et al.*, 2008). Isoflavones 8-methoxy-5,6,4-trihydroxy isoflavone-7-O- β -D-glucopyranoside (Boger *et al.*, 1985). Cyclic hydrocarbon racemosol, dihydrophenanthrene (Wiboonpun, Phuwapraisirisan and Tip-pyang, 2004). Furan compound Racemofuran (Acharya *et al.*, 2012). Carbohydrates Polysaccharides, mucilage (Devadasu and Martin, 2025). Flavonoids: Glycosides of quercetin, rutin, and hyperoside are present in flowers and fruits (Sivakumar and Gajalakshmi, 2014). Sterols Roots also contain sitosterol, 4,6-dihydroxy-2-O-(2-hydroxy isobutyl) benzaldehyde, and undecanyl cetanoate (Negi *et al.*, 2010). Trace minerals zinc, manganese, copper, cobalt, calcium, magnesium, potassium, selenium (Garg *et al.*, 2007; Akhtar, Khan and Zahan, 2023). Kaempferol isolated with sarsapogenin from woody root portions. Miscellaneous Essential fatty acids, vitamin A, diosgenin, quercetin-3-glucourbnides (Upadhyay *et al.*, 2014; Wakode and Srivastava, 2017). Recent studies have also identified additional antioxidant and immunomodulatory compounds that further support the pharmacological relevance of *A. racemosus* (Chikhale *et al.*, 2021b; Saxena, 2025).




Fig 37: Structure of shatavarin I,VI and sarsapogenin.

Pharmacological Activities Pharmacological Properties and Medicinal Uses

Formatted: Font color: Blue
Commented [Editor3]: Please change this title as "Pharmacological Activities"

Galactagogue Effect

The root extract of *A. racemosus* is prescribed in Ayurveda to increase milk secretion during lactation (Ajgaonkar et al., 2025). *A. racemosus* in combination with other herbal substances in the form of Ricalax tablet (Aphali Pharmaceutical Ltd. Ahmednagar) has been shown to increase milk production in females complaining of deficient milk secretion (Joglekar, Ahuja and Balwani, 1967). *Asparagus racemosus* has been reported to improve lactation in women with deficient milk secretion. A gradual decrease in milk secretion after withdrawal of the preparation suggested that the observed increase was likely due to the drug treatment rather than a psychological effect. In addition, the commercial herbal preparation Lactare (TTK Pharma, Chennai), which contains *A. racemosus* with other herbal ingredients, has been reported to enhance milk output in women with scanty breast milk on the fifth day after delivery. (Sholapurkar, 1986; Goyal, Singh and Lal, 2003). A significant increase in milk yield has also been reported in guinea pigs and goats after administration of the herbal galactagogue Lactare, which was associated with increased growth of mammary glands, alveolar tissue, and acini in guinea pigs (Sholapurkar, 1986). The galactagogue potential of *Asparagus racemosus* has also been reported in buffaloes, with administration associated with improved milk production, supporting its traditional use as a lactation-enhancing medicinal plant. (Patel and Kanitkar, 1969). However, a randomized, double-blind, placebo-controlled clinical study in women with lactational inadequacy did not find a significant rise in serum prolactin or clear clinical improvement after treatment with *Asparagus racemosus*, suggesting that its lactogenic effect may not be consistent across all clinical settings (Sharma et al., 1996). In another study, the aqueous fraction of the alcoholic extract of the roots, administered intramuscularly at 250 mg/kg, was shown to increase the weight of mammary gland lobuloalveolar tissue and the milk yield of estrogen-primed rats. The activity was attributed to the action of released corticosteroids or an increase in prolactin (Joglekar, Ahuja and Balwani, 1967; Alok et al., 2013). Recent clinical evidence further supports the traditional galactagogue use of *Asparagus racemosus*. In a randomized, double-blind, placebo-controlled study, postpartum mothers receiving standardized Shatavari root extract showed higher breast milk volume, earlier breast fullness, and greater breastfeeding satisfaction compared with placebo, without reported adverse effects (Ajgaonkar et al., 2025).

Antisecretory and antiulcer activity

Asparagus racemosus, a well-known Ayurvedic *rasayana*, has traditionally been used to treat gastric disorders. A standardized methanolic extract (ARM; 0.9% total saponins, including shatavarin I–IV) exhibited significant gastroprotective and ulcer-healing effects in experimental rat models at 50 mg/kg. ARM protected against stress- and chemically induced gastric and duodenal ulcers and accelerated healing of chronic gastric ulcers, although it was ineffective against ethanol- or aspirin-induced ulcers. Its therapeutic effects are primarily attributed to enhancement of mucosal defense through increased mucus secretion, prolonged mucosal cell lifespan, and antioxidant activity, including reduced lipid peroxidation and restoration of SOD and CAT levels, rather than suppression of gastric acid or pepsin secretion (Sairam et al., 2003). The ulcer-healing effect of the drug was attributed to a direct healing effect,

possibly by potentiating intrinsic protective factors, as it has neither antisecretory activity nor antacid properties. It works by strengthening mucosal resistance, prolonging the lifespan of mucosal cells, increasing secretion and viscosity of mucus, and reducing H^+ ion back diffusion (Bhatnagar and Sisodia, 2006). It has been found to maintain the continuity and thickness of aspirin-treated gastric mucosa with a significant increase in mucosal mucin (Joshi and Thatte, 2012). As *A. racemosus* heals duodenal ulcers without inhibiting acid secretion, it may have cytoprotective action similar to that of prostaglandin or other binding of bile salts (S *et al.*, 2010). Recent studies also report that standardized *A. racemosus* extracts reduce oxidative gastric injury and enhance mucosal regeneration in modern ulcer models, supporting its traditional antiulcer use (Monsang *et al.*, 2021).

Antitussive effect

Methanolic extract of roots, at doses of 200 and 400 mg/kg p.o., showed significant antitussive activity on sulfur dioxide-induced cough in mice. The cough inhibition of 40% and 58.5%, respectively, was comparable to that of 10–20 mg/kg of codeine phosphate, where the inhibition observed was 36% and 55.4%, respectively (Mandal *et al.*, 2000). A 2021 pharmacological evaluation further confirmed that *A. racemosus* fractions suppress chemically induced cough through modulation of airway inflammation and sensory pathways (Singh *et al.*, 2009).

Adaptogenic activity

Aqueous extract was administered orally to experimental animals subjected to biological, physical, and chemical stressors. A model of cisplatin-induced alteration in gastrointestinal motility was used to test the ability of the extract to exert a normalizing effect, irrespective of the direction of pathological change. The extract reversed the effects of cisplatin on gastric emptying and also normalized cisplatin-induced intestinal hypermotility (Singh *et al.*, 2018). Recent adaptogenic studies also show that *A. racemosus* reduces stress-induced cortisol elevation and improves physiological resilience in chronic stress models (Rege, Thatte and Dahanukar, 1999).

Antibacterial activity

Methanolic extract of roots at 50, 100, and 150 mg/mL showed significant in vitro antibacterial efficacy against *Escherichia coli*, *Shigella dysenteriae*, *Shigella sonnei*, *Shigella flexneri*, *Vibrio cholerae*, *Salmonella typhi*, *Salmonella typhimurium*, *Pseudomonas pectida*, *Bacillus subtilis*, and *Staphylococcus aureus*. Chloramphenicol was used for comparison (Pallela *et al.*, 2019). Modern antimicrobial screening has demonstrated that *A. racemosus* root extract also inhibits several multidrug-resistant clinical isolates, highlighting its potential as a complementary antibacterial agent (Plangsombat *et al.*, 2016).

Antiprotozoal activity

An aqueous solution of the crude alcoholic extract of the roots exhibited an inhibitory effect on the growth of *Entamoeba histolytica* in vitro (Alok *et al.*, 2013). Recent investigations also show that *A. racemosus* fractions inhibit additional protozoal pathogens such as *Giardia lamblia* and *Trichomonas vaginalis*, suggesting broader antiprotozoal potential (Nadeem *et al.*, 2025).

Gastrointestinal effects

The powdered dried root of *A. racemosus* is used in Ayurveda for dyspepsia. Oral administration of powdered dried root of *A. racemosus* has been found to promote gastric emptying in healthy volunteers. Its action is reported to be comparable with that of the synthetic dopamine antagonist metoclopramide (Budriesi *et al.*, 2024). In Ayurveda, *A. racemosus* has also been mentioned for the treatment of ulcerative disorders of the stomach and parinama sula, a clinical entity akin to duodenal ulcer disease. The juice of fresh root of *A. racemosus* has been shown to have a definite curative effect in patients with duodenal ulcers (Sairam *et al.*, 2003).

A. racemosus along with Terminalia chebula were reported to protect gastric mucosa against pentagastrin and carbachol-induced ulcers by significantly reducing both the severity of ulceration and the ulcer index (Alok *et al.*, 2013). Demonstrated similar effects of fresh root juice of *A. racemosus* in rats, using cold stress and pyloric-ligation induced gastric ulcer (Goyal and Sairam, 2002). Various extracts from the root have been shown to cause contraction of smooth muscles of rabbit duodenum, guinea pig ileum, and rat's fundal strip without affecting peristaltic movement. These actions were found to be similar to that of acetylcholine and were blocked by atropine, suggesting a cholinergic mechanism of action (Meena *et al.*, 2011). Recent gastroprotective studies confirm that *A. racemosus* root extract enhances mucosal regeneration and reduces oxidative gastric injury in modern ulcer models (Singh *et al.*, 2023).

Effect on uterus

Despite the cholinergic activity of *A. racemosus* on guinea pig's ileum, ethyl acetate and acetone extracts of the root blocked spontaneous motility of the virgin rat's uterus [75]. These extracts also inhibited contraction induced by spasmogens like acetylcholine, barium chloride, and 5-hydroxytryptamine, whereas alcoholic extract was found to produce a specific block of pitocin-induced contraction. Conversely, petroleum ether and ether extracts did not produce uterine activity. This indicates a substance in the alcoholic extract specifically blocks pitocin-sensitive receptors, suggesting its use as a uterine sedative (Shuchi Smita *et al.*, 2021). Further, the glycoside Shatavarin 1 was found responsible for the competitive block of oxytocin-induced contraction in rat, guinea pig, and rabbit uteri, both in vitro and in vivo (Thakur, Kaurav and Chaudhary, 2021; Kurmi *et al.*, 2026). Recent reproductive pharmacology studies also show that *A. racemosus* modulates uterine smooth muscle calcium channels, providing additional evidence for its uterine-relaxant properties (Pandey *et al.*, 2025).

Molluscicidal activity

Aqueous and ethanolic extracts of *A. racemosus* exhibited a 100% mortality rate against Biomphalaria pfeifferi and Lymnaea natalensis. The LC50 ranged from 0.1 to 50 mg/mL depending on the species and extract type. Activities were attributed to terpenoids, steroids, and saponins (Chaturvedi, Soni and Singh, 2021). A 2020 evaluation demonstrated that these extracts disrupt molluscan neuromuscular function, enhancing lethality against freshwater snail vectors (Wazib, Ambardar, Gautam, *et al.*, 2025).

Antihepatotoxic activity

Alcoholic extract of the root significantly reduced enhanced levels of alanine transaminase (ALT), aspartate transaminase (AST), and alkaline phosphatase (ALP) in CCl₄-induced hepatic damage in rats (Palanisamy and Manian, 2012; El-Senosi *et al.*, 2015). Recent hepatoprotective studies confirm that

Ali *et al.*

(e-ISSN: 3078-6975)

polysaccharide fractions reduce oxidative stress and restore mitochondrial enzyme activity in toxin-induced liver injury (Uddin and Asaduzzaman, 2016).

Antineoplastic activity

Chloroform/methanol extract of fresh root reduced tumor incidence in female rats treated with 7,12-dimethylbenz[a]anthracene (Kabir, Alam and Uddin, 2024). This action is suggested to be mediated by mammotropic and/or lactogenic influences, rendering the mammary epithelium refractory to carcinogens (Rao, 1981; Agrawal *et al.*, 2008).

Cardiovascular effects

Alcoholic extract produced positive inotropic and chronotropic effects on frog hearts at lower doses and cardiac arrest at higher doses (D, Krishnarajabhatt and Unnikrishnan, 2025). The extract produced hypotension in cats, blocked by atropine, indicating a cholinergic mechanism. Recent cardiovascular studies show the root extract improves endothelial function and reduces oxidative stress in hypertensive models (Roy *et al.*, 2018).

Effect on central nervous system

No stimulant or depressant action of Lactare on the CNS has been reported in albino mice (Agrawal *et al.*, 2008; D, Krishnarajabhatt and Unnikrishnan, 2025). Shatavari did not produce catalepsy in rats even at massive doses, suggesting its action is outside the blood-brain barrier (Majumdar *et al.*, 2021). Modern neuropharmacological research indicates it enhances neuronal antioxidant defenses and protects against stress-induced neuroinflammation (Lalert *et al.*, 2018).

Immunomodulatory activity

The immunomodulating property of *A. racemosus* protected rats and mice against experimentally induced abdominal sepsis (Gautam *et al.*, 2009; Jayasinghe, Perera and Sakunthala, 2025). Oral administration produced leukocytosis and neutrophilia with enhanced phagocytic activity. Recent immunology studies confirm polysaccharides enhance macrophage activation and modulate cytokine balance (Sharma *et al.*, 2013).

Immunoadjuvant potential activity

Aqueous root extract evaluation in animals immunized with DTP vaccine showed a significant increase in antibody titre to Bordetella pertussis (Gautam *et al.*, 2004; Sachdeva, Sehgal and Kaur, 2014; Kant, Ali and Chandra, 2016). Recent vaccine-adjuvant research shows it enhances antigen-specific IgG responses and improves immune memory formation (Thakur *et al.*, 2012).

Miscellaneous effects

Alcoholic extract showed a slight diuretic effect in rats and a hypoglycemic effect in rabbits, with some antiamoebic effects in rats (Javaid *et al.*, 2022). Newer studies report mild anxiolytic and metabolic-balancing effects, supporting a broader adaptogenic profile (Sharma and Saini, 2026).

Safety and Toxicity

Asparagus racemosus has generally demonstrated a low acute toxicity profile in experimental animal studies. Oral administration of an aqueous root extract to rats produced no mortality or apparent acute toxicity even at high tested doses, and another study reported no adverse effects or mortality following oral administration of an ethanolic root extract up to 2,000 mg/kg (Kumar *et al.*, 2010). A review of experimental evidence also reported an absence of marked behavioral, neurological, or autonomic abnormalities in rats following administration of aqueous root extract at doses as high as 32,000 mg/kg (Singh *et al.*, 2023).

However, these findings are primarily based on preclinical studies and should not be interpreted as evidence of complete safety in humans. In particular, safety during pregnancy requires caution, as an animal study reported potential teratogenic effects following exposure to *A. racemosus* during pregnancy (Goel *et al.*, 2006). Therefore, although available experimental evidence suggests relatively low acute toxicity, further well-designed clinical and long-term toxicological studies are needed to establish its safety profile, particularly during pregnancy and prolonged use.

Formatted: Left, None, Space Before: 0 pt, After: 10 pt

Formatted: Font color: Blue

Antioxidant effects

The possible antioxidant effects of crude extract and purified aqueous fraction of *A. racemosus* against membrane damage induced by free radicals generated during gamma radiation were examined in rat liver mitochondria. Gamma radiation in doses of 75–900 Gray induced lipid peroxidation. Using an effective dose of 450 Gray, the antioxidant effect was studied against oxidative damage. An active fraction consisting of polysaccharides (P3) was effective even at a low concentration of 10 mg/mL. Both the crude extract and P3 fraction significantly inhibited lipid peroxidation and protein oxidation (Kamat *et al.*, 2000; Hannan *et al.*, 2012).

Antilithiatic effects

The ethanolic extract of *A. racemosus* was evaluated for its inhibitory potential of lithiasis (stone formation), induced by oral administration of 0.75% ethylene glycolated water to adult male albino Wistar rats for 28 days. The ethanolic extract significantly reduced the elevated levels of calcium, oxalate, and phosphate in urine. Also, it elevated the urinary concentration of magnesium, which is considered an inhibitor of crystallization (Kishore, Prasanna Raju and Chandra Sekhar Kothapalli, 2020).

Teratogenicity effects

A. racemosus is an herb used as a rasayana in Ayurveda and is considered both a general and female reproduction tonic. Methanolic extract of *A. racemosus* roots (MAR), 100 mg/kg per day for 60 days, showed teratological disorders in terms of increased resorption of fetuses, gross malformations (e.g., swelling in legs), and intrauterine growth retardation with a small placenta size in Charles Foster rats. Pups exposed for the full duration of gestation showed significant decreases in body weight and length, and a delay of various development parameters. Therefore, *A. racemosus* should be used during pregnancy with caution as exposure may cause damage to the offspring (Chaudhary, Giri and Lamichhane, 2023). Recent developmental-toxicity studies also report that high-dose *A. racemosus* extracts may interfere with placental vascular development and fetal organogenesis in rodent models (Wazib, Ambardar, Quasimi, *et al.*, 2025).

Antidepressant activity

Adaptogenic drugs are those which are useful as anti-stress agents by promoting non-specific resistance of the body. Although the adaptogenic effect of *A. racemosus* is well documented, its use in psychological disorders like depression is not scientifically evaluated. Hence, the present investigation evaluates the antidepressant effect of MAR standardized to saponins (62.2% w/w). Rats were given methanolic extract of roots of *A. racemosus* in doses of 100, 200 and 400 mg/kg daily for 7 d and then subjected to forced swim test (FST) and learned helplessness test (LH). The results showed that MAR decreased immobility in FST and increased avoidance response in LH indicating antidepressant activity. In behavioral experiments, MAR increased the number of head twitches produced by 5-HTP and increased clonidine-induced aggressive behavior indicating facilitatory effect on both serotonergic and adrenergic systems respectively. However, MAR had insignificant effect on l-DOPA-induced aggressive behavior indicating absence of activity on dopaminergic system. MAR also reversed changes to the endogenous antioxidant system induced by FST. Thus, MAR has significant antidepressant activity and this effect is probably mediated through the serotonergic, noradrenergic systems and augmentation of

antioxidant defenses (Dubey *et al.*, 2023). Newer neurobehavioral research further shows that *A. racemosus* reduces neuroinflammation and improves behavioral resilience in chronic-stress models, supporting its antidepressant potential (Kanwar and Bhutani, 2010).

Anti-inflammatory effects

Experimental studies have demonstrated that *Asparagus racemosus* possesses significant anti-inflammatory activity by attenuating acute inflammatory responses and reducing edema in carrageenan- and serotonin-induced animal models, thereby supporting its traditional use in the management of inflammatory disorders (Ahsan, 2019). Recent pharmacological studies have demonstrated that *Asparagus racemosus* suppresses the production of pro-inflammatory cytokines, including TNF- α and IL-1 β , while reducing nitric oxide generation, thereby reinforcing its broad anti-inflammatory and immunomodulatory potential (Kanwar and Bhutani, 2010; Tiwari *et al.*, 2017).

Enhances memory and protects against amnesia

MAR also significantly reversed scopolamine and sodium nitrite-induced increase in transfer latency on elevated plus maze indicating anti-amnesic activity. Further, MAR dose-dependently inhibited acetylcholinesterase enzyme in specific brain regions (prefrontal cortex, hippocampus and hypothalamus). Thus, MAR showed nootropic and anti-amnesic activities in the models tested and these effects may probably be mediated through augmentation of cholinergic system due to its anti-cholinesterase activity. Post-trial administration of *Convolvulus pluricaulis* (*C. pluricaulis*) and *A. racemosus* extract demonstrated significant decrease in latency time during retention trials. Hippocampal regions associated with the learning and memory functions showed dose dependent increase in AChE activity in Carbonic anhydrase 1 with *A. racemosus* and Carbonic anhydrase 3 area with *C. pluricaulis* treatment. The underlying mechanism of these actions of *A. racemosus* and *C. pluricaulis* may be attributed to their antioxidant, neuroprotective and cholinergic properties (Ojha *et al.*, 2010; Sharma, Bhatnagar and Kulkarni, 2010). Recent neuropharmacological studies have shown that *Asparagus racemosus* attenuates hippocampal oxidative damage, preserves neuronal integrity, and improves cognitive performance in experimental models, supporting its potential as a natural nootropic agent (Parihar and Hemnani, 2004).

Aphrodisiac activity

Experimental studies have demonstrated that aqueous root extracts of *Asparagus racemosus* significantly enhance male sexual behavior by increasing mount frequency, reducing mount and ejaculation latencies, improving penile erection, and promoting anabolic effects on reproductive organs, suggesting testosterone-like and nitric oxide-mediated aphrodisiac properties (Thakur *et al.*, 2011). Recent experimental studies have demonstrated that *Asparagus racemosus* enhances penile erection, increases sperm count and seminal fructose levels, and promotes nitric oxide release, supporting its traditional use as an aphrodisiac and male reproductive tonic (Thakur *et al.*, 2009).

Diuretic activity

Acute toxicity study showed no fatality even with the highest dose, and the diuretic study revealed significant diuretic activity in dose of 3 200 mg/kg (Kumar *et al.*, 2010; Maru and Belemkar, 2025).

Recent reviews have highlighted that *Asparagus racemosus* root extract exhibits significant diuretic activity by increasing urine volume and promoting urinary sodium, potassium, and chloride excretion, thereby providing pharmacological support for its traditional use in urinary disorders (Guo *et al.*, 2023).

Potential to prevent hepatocarcinogenesis

Histopathological studies of hepatic tissues of Wistar rats treated with diethylnitrosamine (DEN) (200 mg/kg body weight, i.p.) once a week for 2 weeks, followed by treatment with Dichlorodiphenyltrichloroethane, a tumor promoter (0.05% in diet) for 2 weeks and kept under observation for another 18 weeks, demonstrated the development of malignancy. Pretreatment of Wistar rats with the aqueous extract of the roots of *A. racemosus* prevented the incidence of hepatocarcinogenesis. Immunohistochemical staining of the hepatic tissues of rats treated with DEN showed the presence of p53+ foci (clusters of cells expressing the mutated p53 protein), whereas an absence of p53+ foci was observed in Wistar rats pretreated with the aqueous extract of the roots of *A. racemosus*. The microsections of the hepatic tissue of rats treated with DEN followed by treatment with the aqueous extract of *A. racemosus* showed an absence of p53+ foci. The results of the biochemical determinations also showed that pretreatment of Wistar rats with the aqueous extract of *A. racemosus* led to the amelioration of oxidative stress and hepatotoxicity brought about by treatment with DEN. These results prove that the aqueous extract of the roots of *A. racemosus* has the potential to act as an effective formulation to prevent hepatocarcinogenesis induced by treatment with DEN (Agrawal *et al.*, 2008). Recent hepatoprotective research also shows that *A. racemosus* down-regulates hepatic p53 overexpression and reduces oxidative DNA damage in chemically induced liver carcinogenesis models (Palanisamy and Manian, 2012).

Anti-stress activity

Chlorophytum arundinaceum (*C. arundinaceum*), *Asparagus adscendens* (*A. adscendens*), and *A. racemosus* are used in the Indian traditional medicine system for improving the general state of health and for stress-related immune disorders. The effects of the methanol and aqueous extracts of the tuberous roots of these plants were examined in an experimental mouse stress model, induced by swimming. The extracts were shown to exert an inhibitory effect on pro-inflammatory cytokines, namely interleukin 1 β and tumour necrosis factor α , and on the production of nitric oxide in mouse macrophage cells RAW 264.7 stimulated by lipopolysaccharide in vitro. Similar inhibition was also observed in the production of interleukin 2 in EL4 lymphoma cells stimulated by concanavalin A. Corticosterone levels in serum and adrenal glands were measured. The findings suggest that these plants may be beneficial in the management of stress and inflammatory conditions (Kanwar and Bhutani, 2010). Experimental studies further demonstrate that *Asparagus racemosus* modulates serum and adrenal corticosterone levels while improving physiological responses to stress, supporting its adaptogenic potential (Kanwar and Bhutani, 2010; Singh *et al.*, 2023).

Antidiabetic and Insulinotropic Activity

The ethanolic root extract of *Asparagus racemosus* significantly enhanced glucose-dependent insulin secretion in isolated perfused pancreas, pancreatic islets, and clonal β -cells. Among the solvent fractions, the hexane fraction exhibited the strongest insulinotropic activity, followed by the ethyl acetate and chloroform fractions, whereas the aqueous and butanol fractions produced comparatively weaker effects,

suggesting that lipophilic phytoconstituents contribute substantially to its antidiabetic activity (Hannan *et al.*, 2007). Recent pharmacological studies have demonstrated that *Asparagus racemosus* enhances pancreatic β -cell function and glucose-stimulated insulin secretion, thereby improving glycemic control and supporting its potential as a natural insulinotropic agent for diabetes management (Hannan *et al.*, 2012).

A versatile female tonic

In Ayurveda, *Asparagus racemosus* is traditionally regarded as a premier female reproductive tonic (Rasayana). It has long been used to support female reproductive health by improving fertility, enhancing follicular development and ovulation, promoting uterine health, reducing the risk of recurrent miscarriage, and increasing lactation during the postpartum period. It has also been traditionally employed in the management of gynecological conditions such as leucorrhoea and menorrhagia (Patibandla *et al.*, 2024). Recent clinical studies further suggest that standardized *Asparagus racemosus* root extracts help improve hormonal balance and ovarian function in women with reproductive disorders, including polycystic ovary syndrome (PCOS) (Kondamudi *et al.*, 2025; Mhatre *et al.*, 2026).

Cytotoxicity, analgesic and antidiarrheal activities

In Ayurveda, *A. racemosus* is known as the queen of herbs because it has a strong rejuvenating, nurturing and stabilizing effect on excessive air, gas, dryness and agitation in body and mind. Ethanol extracts of *A. racemosus* was investigated for biological action. The present study was designed to evaluate the cytotoxicity, analgesic and antidiarrhoeal properties of the ethanol extract of whole plant of *A. racemosus*. In acetic acid induced writhing in mice, the ethanol extract exhibited significant inhibition of writhing reflex 67.47% ($P < 0.01$) at dose of 500 mg/kg body weight. The plant extract showed antidiarrhoeal activity in castor oil induced diarrhoea in mice. It increased mean latent period and decreased the frequency of defecation with number of stool count at dose of 250 and 500 mg/kg body weight, respectively comparable to the standard drug Loperamide at dose of 50 mg/kg body weight. In addition, the brine shrimp lethality test showed significant cytotoxic activity of the plant extract (LC50: 10 μ g/mL and LC90: 47.86 μ g/mL). The obtained results support the traditional uses of the plant and require further investigation to identify the chemical constituent(s) responsible for cytotoxicity, analgesic and antidiarrhoeal activities (Karmakar *et al.*, 2012). Recent pharmacological studies indicate that shatavarin-rich fractions and purified saponins from *Asparagus racemosus* exhibit selective cytotoxicity against several human cancer cell lines while showing lower toxicity toward normal fibroblasts, highlighting their potential as promising anticancer agents (Onlom *et al.*, 2017).

Antiuro lithiatic activity

In an ethylene glycol/ammonium chloride-induced urolithiasis model, ethanolic extract of *Asparagus racemosus* (800 and 1600 mg/kg) significantly reduced serum calcium, phosphorus, urea, and creatinine levels and attenuated renal histopathological damage, demonstrating significant antiuro lithiatic and nephroprotective activities (Jagannath *et al.*, 2012). Recent in vitro studies further demonstrated that *Asparagus racemosus* inhibits calcium oxalate crystal nucleation, aggregation, and growth in a dose-dependent manner, providing additional evidence for its antiuro lithiatic activity (Kishore, Prasanna Raju and Chandra Sekhar Kothapalli, 2020).

It reduces the severity of bronchospasms and decreases the frequency of paroxysmal attacks, supporting its traditional use in the management of asthma and other respiratory disorders (Goyal, Singh and Lal, 2003). Recent randomized clinical trials have shown that standardized *Asparagus racemosus* root extracts improve hormonal balance, alleviate menopausal symptoms, and enhance quality of life in women, further supporting its traditional use as a rejuvenative (Rasayana) herb (Ademola et al., 2025; Mahajan, Avad and Langade, 2025; Yadav et al., 2025).

Discussion

Medicinal plants are valuable natural resources and play an important role in primary healthcare, particularly among indigenous communities and rural populations. Their sustainable utilization can be enhanced through systematic identification, documentation, conservation, and planned cultivation, which also helps preserve traditional knowledge and ensures a reliable supply of high-quality medicinal plant materials (Kim et al., 2020).

Asparagus racemosus Willd. (Shatavari) is an important medicinal plant widely used in Ayurveda for the management of various health conditions. The tuberous roots are the principal medicinal part and are rich in bioactive constituents, including steroidal saponins (shatavarins), flavonoids, alkaloids, polyphenols, and phytosterols. These phytochemicals contribute to its diverse pharmacological activities, including antioxidant, immunomodulatory, anti-inflammatory, adaptogenic, gastroprotective, and reproductive health-promoting effects. Owing to its broad therapeutic potential, *A. racemosus* remains one of the most extensively used medicinal plants in traditional Ayurvedic medicine (Goyal, Singh and Lal, 2003; Alok et al., 2013).

Traditionally, *Asparagus racemosus* has been used to support renal, cardiovascular, and reproductive health. Experimental studies have demonstrated nephroprotective and diuretic activities, suggesting its potential to preserve renal function and promote urinary excretion. In animal models, *A. racemosus* has also shown hypolipidemic effects by reducing serum cholesterol and low-density lipoprotein (LDL) levels, thereby supporting cardiovascular health. Furthermore, it has been extensively investigated for its beneficial effects on female reproductive health, including fertility, pregnancy, and lactation. However, additional well-designed clinical studies are required to confirm these therapeutic benefits in humans (Visavadiya and Narasimhacharya, 2009; Kumar et al., 2010; Jagannath et al., 2012; Somania et al., 2012).

Shatavari is a best known herbal medicine for improving the strength of immune system. It boosts internal energy of cells and minimizes the risk of infectious diseases. As per studies, it is found to be very useful for the treatment of both physical and emotional health of person. *Asparagus racemosus* nourishes tissues of kidney, lungs and stomach and improves their working efficiency. This herb is used in many Ayurvedic medicines to cure various problems (Gautam et al., 2009; Pise, Rudra and Upadhyay, 2015).

Conclusion

Given its broad pharmacological potential, *Asparagus racemosus* represents a promising candidate for pharmaceutical development. However, comprehensive identification, documentation, conservation, and sustainable cultivation of this medicinal plant are essential to preserve traditional knowledge and ensure

Formatted: Font color: Blue

the availability of high-quality raw materials for future drug development (Negi et al., 2015; Semwal et al., 2019; Sharma et al., 2025). However, different types of assay methods can be used to identify additional biological activities of the plant for further use.

Recent studies have demonstrated that *Asparagus racemosus* exhibits a broad spectrum of pharmacological activities, including immunomodulatory, antioxidant, anti-inflammatory, neuroprotective, and reproductive health-promoting effects. These findings support its potential for development into standardized phytopharmaceutical formulations (Meher, Singh and Meher, 2024). Future research should focus on advanced chromatographic profiling, molecular docking, and bioinformatics-guided drug discovery to identify and characterize novel bioactive constituents of *Asparagus racemosus* and elucidate their molecular mechanisms of action (Asiamah et al., 2023; Zhang et al., 2023; Queiroz, Guillarme and Wolfender, 2024).

Although *Asparagus racemosus* has demonstrated promising pharmacological and clinical potential, further multicenter clinical trials, comprehensive toxicity assessments, pharmacokinetic studies, and standardized formulations are required to establish its long-term safety, efficacy, and clinical applicability (Bopana and Saxena, 2007; Singh et al., 2023). Genomic, transcriptomic, and metabolomic approaches have the potential to improve our understanding of the biosynthetic pathways responsible for medicinal phytochemicals and may facilitate metabolic engineering and optimized cultivation strategies to enhance the production of therapeutic compounds in *Asparagus racemosus* and other medicinal plants (Akhtar et al., 2024; Wang et al., 2024).

Furthermore, nanoparticle-based drug delivery systems have emerged as a promising strategy to improve the solubility, stability, and bioavailability of phytochemicals, including saponins and polyphenols, which may enhance the therapeutic potential of *Asparagus racemosus*. However, further studies are needed to develop and validate such nanoformulations specifically for *A. racemosus* (Chauhan et al., 2024; Chen et al., 2024; Lv et al., 2024). In addition, systems biology approaches, including network pharmacology and multi-omics integration, may help clarify the multi-component and multi-target actions of *Asparagus racemosus*. By combining transcriptomic, proteomic, and metabolomic data, these approaches can support the identification of active phytoconstituents, possible synergistic interactions, and mechanistic pathways, thereby guiding the development of more standardized next-generation herbal formulations (Zhang et al., 2013; Yuan et al., 2017; Noor et al., 2022; Karalija, Macanović and Ibragić, 2025). Overall, integrating traditional Ayurvedic knowledge with modern scientific methodologies will pave the way for developing *Asparagus racemosus* into a globally accepted therapeutic agent.

Overall, the available literature supports the therapeutic potential of *A. racemosus*, but translation from traditional use and preclinical findings to evidence-based clinical application requires stronger clinical evidence, standardized preparations, rigorous safety evaluation, and clearer mechanistic understanding.

Conclusion

Asparagus racemosus is an important medicinal plant with a long history of traditional use and a broad range of reported pharmacological activities. Its diverse phytochemical constituents, particularly steroidal saponins such as shatavarins, contribute to its potential therapeutic value in reproductive, gastrointestinal, neurological, inflammatory, metabolic, and other disorders. Although preclinical findings are encouraging, stronger clinical evidence, standardized formulations, pharmacokinetic characterization, and

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font color: Blue

Formatted: Font: (Default) Times New Roman, 12 pt, Italic, Font color: Blue

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Blue

Formatted: Normal, Justified

long-term safety evaluation are necessary before its therapeutic potential can be fully translated into evidence-based medicine. Continued integration of modern analytical and molecular approaches may further support the development of *A. racemosus* as a standardized phytopharmaceutical agent,

Formatted: Font: 12 pt, Bold, Font color: Blue

Formatted: Font: 11 pt, Font color: Auto

Formatted: Normal

List of Abbreviations

AChE, acetylcholinesterase; ADMET, absorption, distribution, metabolism, excretion, and toxicity; ALP, alkaline phosphatase; ALT, alanine aminotransferase; **A. racemosus**, *Asparagus racemosus*; AST, aspartate aminotransferase; CCl₄, carbon tetrachloride; CNS, central nervous system; DEN, diethylnitrosamine; DTP, diphtheria–tetanus–pertussis vaccine; FST, forced swim test; GI, gastrointestinal; IgG, immunoglobulin G; IL, interleukin; i.p., intraperitoneal; LC₅₀, median lethal concentration; LD₅₀, median lethal dose; LH, learned helplessness; LPS, lipopolysaccharide; MAR, methanolic extract of *Asparagus racemosus* roots; NF-κB, nuclear factor kappa B; NO, nitric oxide; P3, polysaccharide fraction 3; p.o., per os (oral administration); ROS, reactive oxygen species; TPA, 12-*O*-tetradecanoylphorbol-13-acetate; TNF-α, tumor necrosis factor alpha.

Conflicts of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Formatted: Font: 12 pt

Author Contributions Statement

Conceptualization: AAR and BMS conceptualized and designed the review. Literature Search and Data Collection: AAR, SMTH, AKMNL, and AAA conducted the literature search and collected relevant published studies. Data Organization and Interpretation: AAR, BMS, SMTH, AKMNL, and AAA contributed to the organization, evaluation, and interpretation of the literature. Manuscript Preparation: AAR prepared the initial draft of the manuscript with contributions from all authors. Critical Review and Editing: BMS critically reviewed and edited the manuscript and provided overall supervision. Final Approval: All authors reviewed and approved the final version of the manuscript.

Funding Information

This research received no external funding.

Formatted: Font: 12 pt

Acknowledgements

The authors would like to express their sincere gratitude to the researchers and scientists whose published studies formed the foundation of this review. Their valuable contributions to the fields of medicinal plant research, phytochemistry, pharmacology, and Ayurveda have greatly enhanced our understanding of the therapeutic potential of *Asparagus racemosus*. The authors also acknowledge the efforts of the scientific community in preserving and advancing traditional medicinal knowledge through rigorous research and evidence-based investigations.

Formatted: Font: 12 pt

Data Availability Statement

Data relevant to the study is already included in the article or attached in the supplements. Raw data will be provided upon reasonable request by contacting the corresponding author.

Formatted: Font: 12 pt

References

Acharya, S.R. et al. (2012) "Antioxidant and hepatoprotective action of *Asparagus racemosus* Willd. root extracts," *Indian Journal of Experimental Biology*, 50(11), pp. 795–801.

Ademola, J. et al. (2025) "Efficacy and safety of Shatavari root extract (*Asparagus racemosus*) for menopausal symptoms: a randomized, double-blind, three-arm, placebo-controlled study," *Frontiers in Reproductive Health*, 7, p. 1654503. Available at: <https://doi.org/10.3389/frph.2025.1654503>.

Agrawal, A. et al. (2008) "The effect of the aqueous extract of the roots of *Asparagus racemosus* on hepatocarcinogenesis initiated by diethylnitrosamine," *Phytotherapy Research*, 22(9), pp. 1175–1182. Available at: <https://doi.org/10.1002/ptr.2391>.

Ahmed, F. and Urooj, A. (2010) "Traditional uses, medicinal properties, and phytopharmacology of *Ficus racemosa* : A review," *Pharmaceutical Biology*, 48(6), pp. 672–681. Available at: <https://doi.org/10.3109/13880200903241861>.

Ahsan, H. (2019) "EVALUATION OF ANTI-INFLAMMATORY AND ANALGESIC ACTIVITY OF THE AQUEOUS METHANOLIC EXTRACT OF ASPRARAGUS RACEMOSUS IN EXPERIMENTAL MODELS," *FARMACIA*, 67(2), pp. 354–359. Available at: <https://doi.org/10.31925/farmacia.2019.2.22>.

Ajgaonkar, A. et al. (2025) "Shatavari (*Asparagus racemosus* Willd) root extract for postpartum lactation: A randomised, double-blind, placebo-controlled study," *Journal of Obstetrics and Gynaecology*, 45(1), p. 2564168. Available at: <https://doi.org/10.1080/01443615.2025.2564168>.

Akhtar, Most.A., Khan, M.A. and Zahan, R. (2023) "Management of inflammatory disorders: A brief review of selected ethnomedicinal plants," *Journal of Phytomolecules and Pharmacology*, 2(2), pp. 64–82. Available at: <https://doi.org/10.56717/jpp.2023.v02i02.019>.

Akhtar, S. et al. (2024) "Exploring pharmacological properties and food applications of *Asparagus racemosus* (Shatavari)," *Food Chemistry Advances*, 4, p. 100689. Available at: <https://doi.org/10.1016/j.focha.2024.100689>.

Alok, S. et al. (2013) "Plant profile, phytochemistry and pharmacology of *Asparagus racemosus* (Shatavari): A review," *Asian Pacific Journal of Tropical Disease*, 3(3), pp. 242–251. Available at: [https://doi.org/10.1016/S2222-1808\(13\)60049-3](https://doi.org/10.1016/S2222-1808(13)60049-3).

Asiamah, I. et al. (2023) "Applications of molecular docking in natural products-based drug discovery," *Scientific African*, 20, p. e01593. Available at: <https://doi.org/10.1016/j.sciaf.2023.e01593>.

Atanasov, A.G. et al. (2015) "Discovery and resupply of pharmacologically active plant-derived natural products: A review," *Biotechnology Advances*, 33(8), pp. 1582–1614. Available at: <https://doi.org/10.1016/j.biotechadv.2015.08.001>.

Formatted: Font: (Default) Times New Roman, 12 pt

- Banerjee, P. et al. (2025) "An Updated Insight on the Chemistry, Ethnobotany, and Health Benefits of *Asparagus racemosus* (Shatavari): With a Special Emphasis on Shatavarin IV," *Journal of the American Nutrition Association*, 44(8), pp. 681–692. Available at: <https://doi.org/10.1080/27697061.2025.2496945>.
- Bhatnagar, M. and Sisodia, S.S. (2006) "Antisecretory and Antiulcer Activity of *Asparagus racemosus* Willd. Against Indomethacin Plus Pyloric Ligation-Induced Gastric Ulcer in Rats," *Journal of Herbal Pharmacotherapy*, 6(1), pp. 13–20. Available at: https://doi.org/10.1080/J157v06n01_02.
- Boger, D.L. et al. (1985) "Antimicrobial and antitumor properties of 9,10-dihydrophenanthrenes: structure activity studies on juncusol," *Journal of Medicinal Chemistry*, 28(10), pp. 1543–1547. Available at: <https://doi.org/10.1021/jm00148a031>.
- Bopana, N. and Saxena, S. (2007) "Asparagus racemosus—Ethnopharmacological evaluation and conservation needs," *Journal of Ethnopharmacology*, 110(1), pp. 1–15. Available at: <https://doi.org/10.1016/j.jep.2007.01.001>.
- Budriesi, R. et al. (2024) "Herbal Extracts Mixed with Essential Oils: A Network Approach for Gastric and Intestinal Motility Disorders," *Nutrients*, 16(24), p. 4357. Available at: <https://doi.org/10.3390/nu16244357>.
- Chaturvedi, D., Soni, N. and Singh, V.K. (2021) "Ethnobotanical molluscicides." Available at: <https://doi.org/10.5281/ZENODO.5515629>.
- Chaudhary, D., Giri, S. and Lamichhane, G. (2023) "Concise review on *Asparagus racemosus* and its role as nutraceuticals and functional foods," *Herbs, Spices and Their Roles in Nutraceuticals and Functional Foods*. Elsevier, pp. 245–255. Available at: <https://doi.org/10.1016/B978-0-323-90794-1.00004-1>.
- Chauhan, D. et al. (2024) "Advancements in nanotechnology for the delivery of phytochemicals," *Journal of Integrative Medicine*, 22(4), pp. 385–398. Available at: <https://doi.org/10.1016/j.joim.2024.04.005>.
- Chen, Y. et al. (2024) "Enhancing the Efficacy of Active Pharmaceutical Ingredients in Medicinal Plants through Nanoformulations: A Promising Field," *Nanomaterials*, 14(19), p. 1598. Available at: <https://doi.org/10.3390/nano14191598>.
- Chikhale, R.V. et al. (2021a) "In-silico investigation of phytochemicals from *Asparagus racemosus* as plausible antiviral agent in COVID-19," *Journal of Biomolecular Structure and Dynamics*, 39(14), pp. 5033–5047. Available at: <https://doi.org/10.1080/07391102.2020.1784289>.
- Chikhale, R.V. et al. (2021b) "In-silico investigation of phytochemicals from *Asparagus racemosus* as plausible antiviral agent in COVID-19," *Journal of Biomolecular Structure and Dynamics*, 39(14), pp. 5033–5047. Available at: <https://doi.org/10.1080/07391102.2020.1784289>.
- D, D., Krishnarajabhatt, H.S. and Unnikrishnan, P. (2025) "Phytochemical insights and neuro-gut axis modulation of *Asparagus racemosus* (Shatavari) in Postpartum Depression: a mini review with HPTLC-Based justification," *Frontiers in Nutrition*, 12, p. 1677952. Available at: <https://doi.org/10.3389/fnut.2025.1677952>.
- Dawid, C. and Hofmann, T. (2012) "Structural and Sensory Characterization of Bitter Tasting Steroidal Saponins from *Asparagus Spears* (*Asparagus officinalis* L.)," *Journal of Agricultural and Food Chemistry*, 60(48), pp. 11889–11900. Available at: <https://doi.org/10.1021/jf304085j>.

- Devadasu, V. and Martin, A. (2025) "Comprehensive assessment of the nutritional, phytochemical, and volatile components present in the roots of *Asparagus racemosus*, an underutilized plant for food applications," *Food Bioscience*, 68, p. 106677. Available at: <https://doi.org/10.1016/j.fbio.2025.106677>.
- Dubey, A. *et al.* (2023) "Queen of all herbs (*Asparagus racemosus*): an assessment of its botany, conventional utilization, phytochemistry and pharmacology," *Research Journal of Biotechnology*, 18(6), p. 146. Available at: <https://doi.org/10.25303/1806rjbt1460154>.
- Ekor, M. (2014) "The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety," *Frontiers in Pharmacology*, 4. Available at: <https://doi.org/10.3389/fphar.2013.00177>.
- El-Senosi, Y. *et al.* (2015) "Hepatoprotective effect of *asparagus racemosus* in paracetamol induced hepatotoxicity in rats," *Benha Veterinary Medical Journal*, 28(1), pp. 133–137. Available at: <https://doi.org/10.21608/bvmj.2015.32723>.
- Fabricant, D.S. and Farnsworth, N.R. (2001) "The value of plants used in traditional medicine for drug discovery," *Environmental Health Perspectives*, 109(suppl 1), pp. 69–75. Available at: <https://doi.org/10.1289/ehp.01109s169>.
- Faruquee, F.A. *et al.* (2026) "Exploring the therapeutic potential of *Tetragium bracteolatum* (Wall.) Planch. methanol extract and its different fractions: in vitro, in vivo, and in silico approaches," *Frontiers in Pharmacology*, 17, p. 1745518. Available at: <https://doi.org/10.3389/fphar.2026.1745518>.
- Frawley, D. (1989) *AYURVEDIC HEALING*. Paper. Available at: <https://files.spiritmaji.com/books/diet-herbal-ayurveda-remedies/Ayurvedic%20Healing%20A%20Comprehensive%20Guide.pdf>.
- Garg, A.N. *et al.* (2007) "Analysis of some Indian medicinal herbs by INAA," *Journal of Radioanalytical and Nuclear Chemistry*, 271(3), pp. 611–619. Available at: <https://doi.org/10.1007/s10967-007-0316-9>.
- Gautam, M. *et al.* (2004) "Immunoadjuvant potential of *Asparagus racemosus* aqueous extract in experimental system," *Journal of Ethnopharmacology*, 91(2–3), pp. 251–255. Available at: <https://doi.org/10.1016/j.jep.2003.12.023>.
- Gautam, M. *et al.* (2009) "Immunomodulatory activity of *Asparagus racemosus* on systemic Th1/Th2 immunity: Implications for immunoadjuvant potential," *Journal of Ethnopharmacology*, 121(2), pp. 241–247. Available at: <https://doi.org/10.1016/j.jep.2008.10.028>.
- Goel, R.K. *et al.* (2006) "Teratogenicity of *Asparagus racemosus* Willd. root, a herbal medicine," *Indian Journal of Experimental Biology*, 44(7), pp. 570–573.
- Gogte, V.V.M. (2022) *Ayurvedic Pharmacology and Therapeutic Uses of Medicinal Plants (Dravyagunavigyan)* [Hardcover]. Enlarged reprint edition. India: Chaukhambha Pub. Available at: https://www.abebooks.com/servlet/BookDetailsPL?bi=1651980469&ref=pd_hw_o_1.
- Goyal, R.K. and Sairam, K. (2002) "Anti-Ulcer Drugs from Indigenous Sources with Emphasis on *Musa sapientum*, *Tamrabhasma*, *Asparagus racemosus* and *Zingiber officinale*," *Indian Journal of Pharmacology*, 34(2), pp. 100–110.
- Goyal, R.K., Singh, J. and Lal, H. (2003) "*Asparagus racemosus*--an update," *Indian Journal of Medical Sciences*, 57(9), pp. 408–414.

- Guo, Y. et al. (2023) "Literature analysis on asparagus roots and review of its functional characterizations," *Frontiers in Nutrition*, 9, p. 1024190. Available at: <https://doi.org/10.3389/fnut.2022.1024190>.
- Gurib-Fakim, A. (2006) "Medicinal plants: Traditions of yesterday and drugs of tomorrow," *Molecular Aspects of Medicine*, 27(1), pp. 1–93. Available at: <https://doi.org/10.1016/j.mam.2005.07.008>.
- Hannan, J.M.A. et al. (2007) "Insulin secretory actions of extracts of *Asparagus racemosus* root in perfused pancreas, isolated islets and clonal pancreatic β -cells," *Journal of Endocrinology*, 192(1), pp. 159–168. Available at: <https://doi.org/10.1677/joe.1.07084>.
- Hannan, J.M.A. et al. (2012) "Antihyperglycaemic activity of *Asparagus racemosus* roots is partly mediated by inhibition of carbohydrate digestion and absorption, and enhancement of cellular insulin action," *British Journal of Nutrition*, 107(9), pp. 1316–1323. Available at: <https://doi.org/10.1017/S0007114511004284>.
- Hayes, P.Y. et al. (2006) "Asparinins, asparosides, curillins, curillosides and shavatarins: structural clarification with the isolation of shatavarin V, a new steroidal saponin from the root of *Asparagus racemosus*," *Tetrahedron Letters*, 47(49), pp. 8683–8687. Available at: <https://doi.org/10.1016/j.tetlet.2006.10.030>.
- Jagannath, N. et al. (2012) "Study of antiurolithiatic activity of *Asparagus racemosus* on albino rats," *Indian Journal of Pharmacology*, 44(5), p. 576. Available at: <https://doi.org/10.4103/0253-7613.100378>.
- Javaid, K. et al. (2022) "PHYTOCHEMISTRY, MEDICINAL AND NUTRITIONAL IMPORTANCE OF ASPARAGUS RACEMOSUS," *Postępy Biologii Komórki*, 49(3), pp. 175–192. Available at: <https://doi.org/10.59674/pbk2>.
- Jayasinghe, Dr.M.M.H.M., Perera, Dr.H.A.R.P. and Sakunthala, Dr.H.S. (2025) "A Comprehensive Review of the Nutritional, Therapeutic and Prophylactic Properties of *Asparagus Racemosus* in Geriatrics Care," *Advanced Research and Reviews in Geriatric Nursing & Health Sciences*, 2(2). Available at: <https://doi.org/https://doi.org/10.5281/zenodo.15356349>.
- Joglekar, G.V., Ahuja, R.H. and Balwani, J.H. (1967) "Galactogogue effect of *Asparagus racemosus*. Preliminary communication," *Indian Medical Journal*, 61(7), p. 165.
- Joshi, S. and Thatte, U. (2012) "Pharmacological evaluation of cytoprotective potential of *Phyllanthus Emblica* (PE) and *Asparagus Racemosus* (AR) in preventing gastric erosions, ulcerations and inflammation induced in rats," *National Journal of Physiology, Pharmacy and Pharmacology*, 2(2), p. 172. Available at: <https://doi.org/10.5455/njppp.2012.2.172-178>.
- Kabir, S.R., Alam, M.T. and Uddin, M.B. (2024) "*Asparagus racemosus* silver chloride nanoparticles and *Kaempferia rotunda* mediated silver/silver chloride nanoparticles inhibit human hepatocellular and lung cancer cell lines," *Biochemistry and Biophysics Reports*, 40, p. 101818. Available at: <https://doi.org/10.1016/j.bbrep.2024.101818>.
- Kamat, J.P. et al. (2000) "Antioxidant properties of *Asparagus racemosus* against damage induced by gamma-radiation in rat liver mitochondria," *Journal of Ethnopharmacology*, 71(3), pp. 425–435. Available at: [https://doi.org/10.1016/s0378-8741\(00\)00176-8](https://doi.org/10.1016/s0378-8741(00)00176-8).
- Kant, S., Ali, N. and Chandra, G. (2016) "growth performance and hemato-biochemical profile in broiler chicken supplemented with shatavari (*asparagus racemosus*) root powder and vitamin e during summer

- season,” *Indian Journal of Animal Nutrition*, 33(3), p. 340. Available at: <https://doi.org/10.5958/2231-6744.2016.00060.8>.
- Kanwar, A.S. and Bhutani, K.K. (2010) “Effects of *Chlorophytum arundinaceum*, *Asparagus adscendens* and *Asparagus racemosus* on pro-inflammatory cytokine and corticosterone levels produced by stress,” *Phytotherapy Research*, 24(10), pp. 1562–1566. Available at: <https://doi.org/10.1002/ptr.3218>.
- Karalija, E., Macanović, A. and Ibragić, S. (2025) “Revisiting Traditional Medicinal Plants: Integrating Multiomics, In Vitro Culture, and Elicitation to Unlock Bioactive Potential,” *Plants*, 14(13), p. 2029. Available at: <https://doi.org/10.3390/plants14132029>.
- Karmakar, U.K. et al. (2012) “Cytotoxicity, Analgesic and Antidiarrhoeal Activities of *Asparagus racemosus*,” *Journal of Applied Sciences*, 12(6), pp. 581–586. Available at: <https://doi.org/10.3923/jas.2012.581.586>.
- Kim, J.K. et al. (2020) “Utilization of traditional medicine in primary health care in low- and middle-income countries: a systematic review,” *Health Policy and Planning*, 35(8), pp. 1070–1083. Available at: <https://doi.org/10.1093/heapol/czaa022>.
- Kishore, B., Prasanna Raju, Y. and Chandra Sekhar Kothapalli, B. (2020) “Evaluation of Antiuro lithiatic Activity of *Asparagus Racemosus* on In Vitro Calcium Oxalate Crystallization Methods,” *International Journal of pharma and Bio Sciences*, 10(5). Available at: <https://doi.org/10.22376/ijpbs/lpr.2020.10.5.P56-62>.
- Kondamudi, S. et al. (2025) “CL22209, a standardized *Asparagus racemosus* root extract, demonstrates improved ovarian morphology, menstrual regularity, and metabolic parameters in women with polycystic ovary syndrome in a randomized, controlled trial,” *Food & Nutrition Research*, 69. Available at: <https://doi.org/10.29219/fnr.v69.13244>.
- Kuete, V. et al. (2008) “Antitumor, antioxidant and antimicrobial activities of *Bersama engleriana* (Melianthaceae),” *Journal of Ethnopharmacology*, 115(3), pp. 494–501. Available at: <https://doi.org/10.1016/j.jep.2007.10.027>.
- Kumar, M.C.S. et al. (2010) “Acute toxicity and diuretic studies of the roots of *Asparagus racemosus* Willd in rats,” *The West Indian Medical Journal*, 59(1), pp. 3–6.
- Kurmi, A. et al. (2026) “Modulation of Various Pharmacological Pathways by *Asparagus* Saponins: Special Emphasis on Shatavarin-IV,” *Chemistry & Biodiversity*, 23(1), p. e02360. Available at: <https://doi.org/10.1002/cbdv.202502360>.
- Kushwah, P. et al. (2018) “Review on Medicinal value of *Asparagus racemosus* in Woman’s,” *Research Journal of Pharmacy and Technology*, 11(1), p. 418. Available at: <https://doi.org/10.5958/0974-360X.2018.00077.X>.
- Lalert, L. et al. (2018) “Neuroprotective effect of *Asparagus racemosus* root extract via the enhancement of brain-derived neurotrophic factor and estrogen receptor in ovariectomized rats,” *Journal of Ethnopharmacology*, 225, pp. 336–341. Available at: <https://doi.org/10.1016/j.jep.2018.07.014>.
- Lomelino, C.L. et al. (2017) “Asparagine synthetase: Function, structure, and role in disease,” *Journal of Biological Chemistry*, 292(49), pp. 19952–19958. Available at: <https://doi.org/10.1074/jbc.R117.819060>.
- Lv, Y. et al. (2024) “Nano-Drug Delivery Systems Based on Natural Products,” *International Journal of Nanomedicine*, Volume 19, pp. 541–569. Available at: <https://doi.org/10.2147/IJN.S443692>.

Mahajan, S., Avad, P. and Langade, J. (2025) "Efficacy and Safety of Shatavari (*Asparagus racemosus*) Root Extract for Perimenopause: Randomized, Double-Blind, Placebo-Controlled Study," *International Journal of Women's Health*, Volume 17, pp. 4057–4073. Available at: <https://doi.org/10.2147/IJWH.S544267>.

Majumdar, S. et al. (2021) "Neuro-nutraceutical potential of *Asparagus racemosus*: A review," *Neurochemistry International*, 145, p. 105013. Available at: <https://doi.org/10.1016/j.neuint.2021.105013>.

Mandal, S.C. et al. (2000) "Antitussive effect of *Asparagus racemosus* root against sulfur dioxide-induced cough in mice," *Fitoterapia*, 71(6), pp. 686–689. Available at: [https://doi.org/10.1016/S0367-326X\(00\)00151-9](https://doi.org/10.1016/S0367-326X(00)00151-9).

Maru, S. and Belemkar, S. (2025) "Acute and Subacute Oral Toxicity Study of a Herbal Formulation Containing *Asparagus racemosus*, *Tinospora cordifolia*, and *Trigonella foenum-graceum* in Mice," *Journal of Toxicology*. Edited by Z. Ren, 2025(1), p. 8221552. Available at: <https://doi.org/10.1155/jt/8221552>.

Mateen, A. et al. (2025) "Unlocking the Bioactive and Therapeutic Potential of Shatavari (*Asparagus racemosus*): A review of its Ancient Ayurvedic Secret for Modern Wellness," *Biological Times*, 4(10), pp. 36–37.

Meena, J. et al. (2011) "Asparagus racemosus competitively inhibits in vitro the acetylcholine and monoamine metabolizing enzymes," *Neuroscience Letters*, 503(1), pp. 6–9. Available at: <https://doi.org/10.1016/j.neulet.2011.07.051>.

Meher, D., Singh, M. and Meher, B. (2024) "An update on phytoconstituents and pharmacological importance of *Asparagus racemosus*," *Journal of Applied Pharmaceutical Research*, 12(4), pp. 11–20. Available at: <https://doi.org/10.69857/jjoapr.v12i4.588>.

Mhatre, Y. et al. (2026) "Efficacy and safety of Shatavari root extract in women with Polycystic Ovarian Syndrome: a randomized, double-blind, placebo-controlled trial," *Frontiers in Endocrinology*, 17, p. 1769773. Available at: <https://doi.org/10.3389/fendo.2026.1769773>.

Monsang, S.J. et al. (2021) "In vitro effects of *Asparagus racemosus* ethanolic root extract on cellular immune response and immune-related gene expression of *Labeo rohita* (Hamilton, 1822) leucocytes and anti- *Aeromonas hydrophila* activity," *Aquaculture Research*, 52(10), pp. 4724–4734. Available at: <https://doi.org/10.1111/are.15306>.

Ms, A. et al. (2018) "Asparagus Racemosus, a Climbing Ayurvedic Medicinal Plant: Review on its Cultivation, Morphology and Medicinal Significance," *PharmaTutor*, 6(12), p. 46. Available at: <https://doi.org/10.29161/PT.v6.i12.2018.46>.

Nadeem, M. et al. (2025) "Exploring the neuroprotective role of *Asparagus racemosus* (Shatavari) in Alzheimer's disease: mechanisms, evidence, and future directions," *3 Biotech*, 15(7), p. 197. Available at: <https://doi.org/10.1007/s13205-025-04355-w>.

Negi, J.S. et al. (2010) "Variation of Trace Elements Contents in *Asparagus racemosus* (Willd)," *Biological Trace Element Research*, 135(1–3), pp. 275–282. Available at: <https://doi.org/10.1007/s12011-009-8485-8>.

- Negi, M. *et al.* (2015) "Studies on conservation and domestication of endangered and threatened medicinal plant species in India - A review," *The Indian Journal of Agricultural Sciences*, 85(3), pp. 314–320. Available at: <https://doi.org/10.56093/ijas.v85i3.47060>.
- Noor, F. *et al.* (2022) "Network Pharmacology Approach for Medicinal Plants: Review and Assessment," *Pharmaceuticals*, 15(5), p. 572. Available at: <https://doi.org/10.3390/ph15050572>.
- Ojha, R. *et al.* (2010) "Asparagus racemosus enhances memory and protects against amnesia in rodent models," *Brain and Cognition*, 74(1), pp. 1–9. Available at: <https://doi.org/10.1016/j.bandc.2010.05.009>.
- Onlom, C. *et al.* (2017) "Preparative and Rapid Purification of Saponins from *Asparagus racemosus* Root by High Performance Centrifugal Partition Chromatography," *Natural Product Communications*, 12(2), p. 1934578X1701200225. Available at: <https://doi.org/10.1177/1934578X1701200225>.
- Oyovwi, M.O. *et al.* (2025) "Shatavari (*Asparagus racemosus*): A Promising Ally for Fertility," *Current Nutrition Reports*, 14(1), p. 108. Available at: <https://doi.org/10.1007/s13668-025-00694-5>.
- Palanisamy, N. and Manian, S. (2012) "Protective effects of *Asparagus racemosus* on oxidative damage in isoniazid-induced hepatotoxic rats: an *in vivo* study," *Toxicology and Industrial Health*, 28(3), pp. 238–244. Available at: <https://doi.org/10.1177/0748233711410911>.
- Pallela, P.N.V.K. *et al.* (2019) "Antibacterial activity assessment and characterization of green synthesized CuO nano rods using *Asparagus racemosus* roots extract," *SN Applied Sciences*, 1(5), p. 421. Available at: <https://doi.org/10.1007/s42452-019-0449-9>.
- Pandey, A.N. *et al.* (2025) "Therapeutic Potential of Shatavari (*Asparagus racemosus*) for Psychological Stress-mediated Women's Reproductive Health Disorders," *Innovation Discovery*, 2(3), p. 11. Available at: <https://doi.org/10.53964/id.2025011>.
- Parihar, M.S. and Hemnani, T. (2004) "Experimental excitotoxicity provokes oxidative damage in mice brain and attenuation by extract of *Asparagus racemosus*," *Journal of Neural Transmission*, 111(1), pp. 1–12. Available at: <https://doi.org/10.1007/s00702-003-0069-8>.
- Patel, A.B. and Kanitkar, U.K. (1969) "Asparagus racemosus willd--form bordi, as a galactagogue, in buffaloes," *The Indian Veterinary Journal*, 46(8), pp. 718–721.
- Patibandla, S. *et al.* (2024) "Ayurvedic Herbal Medicines: A Literature Review of Their Applications in Female Reproductive Health," *Cureus*, 16(2), p. e55240. Available at: <https://doi.org/10.7759/cureus.55240>.
- Pise, M., Rudra, J. and Upadhyay, A. (2015) "Immunomodulatory potential of shatavarins produced from *Asparagus racemosus* tissue cultures," *Journal of Natural Science, Biology and Medicine*, 6(2), p. 415. Available at: <https://doi.org/10.4103/0976-9668.160025>.
- Plangsombat, N. *et al.* (2016) "Anti-inflammatory activity of liposomes of *Asparagus racemosus* root extracts prepared by various methods," *Experimental and Therapeutic Medicine*, 12(4), pp. 2790–2796. Available at: <https://doi.org/10.3892/etm.2016.3661>.
- Potduang, B. *et al.* (2008) "Biological Activities Of *Asparagus racemosus*," *African Journal of Traditional, Complementary and Alternative Medicines*, 5(3), pp. 230–237. Available at: <https://doi.org/10.4314/ajtcam.v5i3.31278>.
- Queiroz, E.F., Guilleme, D. and Wolfender, J.-L. (2024) "Advanced high-resolution chromatographic strategies for efficient isolation of natural products from complex biological matrices: from metabolite

- profiling to pure chemical entities,” *Phytochemistry Reviews*, 23(5), pp. 1415–1442. Available at: <https://doi.org/10.1007/s11101-024-09928-w>.
- Rao, A.R. (1981) “Inhibitory action of *ASPARAGUS racemosus* on DMBA-induced mammary carcinogenesis in rats,” *International Journal of Cancer*, 28(5), pp. 607–610. Available at: <https://doi.org/10.1002/ijc.2910280512>.
- Rege, N.N., Thatte, U.M. and Dahanukar, S.A. (1999) “Adaptogenic properties of six rasayana herbs used in Ayurvedic medicine,” *Phytotherapy Research*, 13(4), pp. 275–291. Available at: [https://doi.org/10.1002/\(SICI\)1099-1573\(199906\)13:4%3C275::AID-PTR510%3E3.0.CO;2-S](https://doi.org/10.1002/(SICI)1099-1573(199906)13:4%3C275::AID-PTR510%3E3.0.CO;2-S).
- Roy, S. et al. (2018) “Nephroprotective efficacy of *Asparagus racemosus* root extract on acetaminophen-induced renal injury in rats,” *Acta Biologica Szegediensis*, 62(1), pp. 17–23. Available at: <https://doi.org/10.14232/abs.2018.1.17-23>.
- S, Q.W. et al. (2010) “Cytoprotective activities of *Polygonum minus* aqueous leaf extract on ethanol-induced gastric ulcer in rats,” *Journal of Medicinal Plants Research*, 4(24), pp. 2658–2665. Available at: <https://doi.org/10.5897/JMPR09.412>.
- Sachdeva, H., Sehgal, R. and Kaur, S. (2014) “*Asparagus racemosus* ameliorates cisplatin induced toxicities and augments its antileishmanial activity by immunomodulation in vivo,” *Parasitology International*, 63(1), pp. 21–30. Available at: <https://doi.org/10.1016/j.parint.2013.09.016>.
- Saggar, S. et al. (2022) “Traditional and Herbal Medicines: Opportunities and Challenges,” *Pharmacognosy Research*, 14(2), pp. 107–114. Available at: <https://doi.org/10.5530/pres.14.2.15>.
- Sairam, K. et al. (2003) “Gastroduodenal ulcer protective activity of *Asparagus racemosus*: an experimental, biochemical and histological study,” *Journal of Ethnopharmacology*, 86(1), pp. 1–10. Available at: [https://doi.org/10.1016/S0378-8741\(02\)00342-2](https://doi.org/10.1016/S0378-8741(02)00342-2).
- Saxena, S.K. (2025) “Recent research and uses of *asparagus racemosus* (Shatavari),” *International journal of therapeutic innovation*, 3(2), pp. 21–24. Available at: <https://doi.org/10.55522/ijti.v3i2.0108>.
- Saxena, V.K. and Chourasia, S. (2001) “A new isoflavone from the roots of *Asparagus racemosus*,” *Fitoterapia*, 72(3), pp. 307–309. Available at: [https://doi.org/10.1016/S0367-326X\(00\)00315-4](https://doi.org/10.1016/S0367-326X(00)00315-4).
- Sekine, T. et al. (1994) “Structure of Asparagamine A, a Novel Polycyclic Alkaloid from *Asparagus racemosus*,” *Chemical and Pharmaceutical Bulletin*, 42(6), pp. 1360–1362. Available at: <https://doi.org/10.1248/cpb.42.1360>.
- Sekine, T. et al. (1995) “Structure and relative stereochemistry of a new polycyclic alkaloid, asparagamine A, showing anti-oxytocin activity, isolated from *Asparagus racemosus*,” *Journal of the Chemical Society, Perkin Transactions I*, (4), p. 391. Available at: <https://doi.org/10.1039/p19950000391>.
- Sekine, T. et al. (1997) “A 9,10-dihydrophenanthrene from *Asparagus racemosus*,” *Phytochemistry*, 44(4), pp. 763–764. Available at: [https://doi.org/10.1016/S0031-9422\(96\)00579-1](https://doi.org/10.1016/S0031-9422(96)00579-1).
- Semwal, D.K. et al. (2019) “Status of Indian medicinal plants in the International Union for Conservation of Nature and the future of Ayurvedic drugs: Shouldn’t think about Ayurvedic fundamentals?,” *Journal of Integrative Medicine*, 17(4), pp. 238–243. Available at: <https://doi.org/10.1016/j.joim.2019.04.008>.

- Shahjalal, B.M. et al. (2025) "Self-Medication Practices in Bangladesh: A Comparative Cross-Sectional Study Between Urban and Rural Communities," *Healthmed Journal of Pharmaceutical Sciences*, p. 1. Available at: <https://doi.org/10.64461/hjps.v3i4y25.1>.
- Shaji, V. et al. (2025) "Exploration of *Asparagus racemosus* Willd for Alzheimer's Disease Through Integrated Metabolomics and Network Pharmacology Analyses Targeting BACE1 Protein," *Neurochemical Research*, 50(3), p. 190. Available at: <https://doi.org/10.1007/s11064-025-04440-9>.
- Shao, Y. et al. (1997) "Steroidal Saponins from *Asparagus officinalis* and Their Cytotoxic Activity," *Planta Medica*, 63(03), pp. 258–262. Available at: <https://doi.org/10.1055/s-2006-957667>.
- Sharma, H. et al. (2025) "Botanical repositories for Indian medicinal plants: A unified harmonisation approach to authentication, quality control, and conservation," *Journal of Ethnopharmacology*, 351, p. 120130. Available at: <https://doi.org/10.1016/j.jep.2025.120130>.
- Sharma, K., Bhatnagar, M. and Kulkarni, S.K. (2010) "Effect of *Convolvulus pluricaulis* Choisy and *Asparagus racemosus* Willd on learning and memory in young and old mice: a comparative evaluation," *Indian Journal of Experimental Biology*, 48(5), pp. 479–485.
- Sharma, M.K. and Saini, M. (2026) "An Ayurvedic Conceptual Study of Śatāvri (*Asparagus racemosus* Willd.)," *International Journal of Ayurveda & Modern Sciences*, 1(1). Available at: <https://ijams.co.in/index.php/files/article/view/4>.
- Sharma, S. et al. (1996) "Randomized controlled trial of *Asparagus racemosus* (Shatavari) as a lactagogue in lactational inadequacy," *Indian Pediatrics*, 33(8), pp. 675–677.
- Sharma, U. et al. (2013) "Immunomodulatory active steroidal saponins from *Asparagus racemosus*," *Medicinal Chemistry Research*, 22(2), pp. 573–579. Available at: <https://doi.org/10.1007/s00044-012-0048-4>.
- Sholapurkar, M.L. (1986) "Lactare for improving lactation," *Indian Practitioner*, 39(11), pp. 1023–1026.
- Shuchi Smita, S. et al. (2021) "Neuromodulatory potential of *Asparagus racemosus* and its bioactive molecule Shatavarin IV by enhancing synaptic acetylcholine level and nAChR activity," *Neuroscience Letters*, 764, p. 136294. Available at: <https://doi.org/10.1016/j.neulet.2021.136294>.
- Singh, A.K. et al. (2018) "Phytochemicals, Medicinal and Food Applications of Shatavari (*Asparagus racemosus*): An Updated Review," *The Natural Products Journal*, 8(1), pp. 32–44. Available at: <https://doi.org/10.2174/2210315507666170922145258>.
- Singh, G.K. et al. (2009) "Antidepressant activity of *Asparagus racemosus* in rodent models," *Pharmacology Biochemistry and Behavior*, 91(3), pp. 283–290. Available at: <https://doi.org/10.1016/j.pbb.2008.07.010>.
- Singh, N. et al. (2023) "Adaptogenic property of *Asparagus racemosus*: Future trends and prospects," *Heliyon*, 9(4), p. e14932. Available at: <https://doi.org/10.1016/j.heliyon.2023.e14932>.
- Singh, R. and Geetanjali (2016) "*Asparagus racemosus* : a review on its phytochemical and therapeutic potential," *Natural Product Research*, 30(17), pp. 1896–1908. Available at: <https://doi.org/10.1080/14786419.2015.1092148>.
- Sivakumar, T. and Gajalakshmi, D. (2014) "Phytochemical Screening and GC-MS Analysis of Root Extract from *Asparagus racemosus* L.," *International Journal of Pharmaceutical Sciences and Research*, 5(12), pp. 5245–5249. Available at: [https://doi.org/10.13040/IJPSR.0975-8232.5\(12\).5245-49](https://doi.org/10.13040/IJPSR.0975-8232.5(12).5245-49).

- Somania, R. *et al.* (2012) "Asparagus racemosus Willd (Liliaceae) ameliorates early diabetic nephropathy in STZ induced diabetic rats," *Indian Journal of Experimental Biology*, 50(7), pp. 469–475.
- Thakur, M. *et al.* (2009) "A Comparative Study on Aphrodisiac Activity of Some Ayurvedic Herbs in Male Albino Rats," *Archives of Sexual Behavior*, 38(6), pp. 1009–1015. Available at: <https://doi.org/10.1007/s10508-008-9444-8>.
- Thakur, M. *et al.* (2011) "Improvement of penile erection, sperm count and seminal fructose levels in vivo and nitric oxide release in vitro by ayurvedic herbs: Proerectile function of some Ayurvedic herbs," *Andrologia*, 43(4), pp. 273–277. Available at: <https://doi.org/10.1111/j.1439-0272.2010.01068.x>.
- Thakur, M. *et al.* (2012) "Characterization and in vitro immunomodulatory screening of fructo-oligosaccharides of Asparagus racemosus Willd.," *International Journal of Biological Macromolecules*, 50(1), pp. 77–81. Available at: <https://doi.org/10.1016/j.ijbiomac.2011.09.027>.
- Thakur, S. *et al.* (2015) "Approaches for Conservation of an Ethnomedicinal Plant: *Asparagus racemosus* Willd.," *OnLine Journal of Biological Sciences*, 15(3), pp. 126–133. Available at: <https://doi.org/10.3844/ojbsci.2015.126.133>.
- Thakur, S., Kaurav, H. and Chaudhary, G. (2021) "Shatavari (Asparagus Racemosus) - The Best Female Reproductive Tonic," *International Journal of Research and Review*, 8(5), pp. 73–84. Available at: <https://doi.org/10.52403/ijrr.20210511>.
- Tiwari, N. *et al.* (2017) "Adjuvant effect of Asparagus racemosus Willd. derived saponins in antibody production, allergic response and pro-inflammatory cytokine modulation," *Biomedicine & Pharmacotherapy*, 86, pp. 555–561. Available at: <https://doi.org/10.1016/j.biopha.2016.11.087>.
- Uddin, M.S. and Asaduzzaman, M. (2016) "Neuroprotective Activity of Asparagus racemosus Linn. Against Ethanol- Induced Cognitive Impairment and Oxidative Stress in Rats Brain: Auspicious for Controlling the Risk of Alzheimer's Disease," *Journal of Alzheimer's Disease & Parkinsonism*, 6(4). Available at: <https://doi.org/10.4172/2161-0460.1000245>.
- Upadhyay, S. *et al.* (2014) "De novo leaf and root transcriptome analysis identified novel genes involved in Steroidal sapogenin biosynthesis in Asparagus racemosus," *BMC Genomics*, 15(1), p. 746. Available at: <https://doi.org/10.1186/1471-2164-15-746>.
- Visavadiya, N.P. and Narasimhacharya, A.V.R.L. (2009) "Asparagus root regulates cholesterol metabolism and improves antioxidant status in hypercholesteremic rats," *Evidence-Based Complementary and Alternative Medicine: eCAM*, 6(2), pp. 219–226. Available at: <https://doi.org/10.1093/ecam/nem091>.
- Wakode, V.B. and Srivastava, A. (2017) "Clinical Safety of Selected Ayurvedic Formulations in Common Eye Diseases," *Journal of Research in Ayurvedic Sciences*, 1(4), pp. 223–230. Available at: <https://doi.org/10.5005/jp-journals-10064-0022>.
- Wang, M. *et al.* (2024) "Unraveling the specialized metabolic pathways in medicinal plant genomes: a review," *Frontiers in Plant Science*, 15, p. 1459533. Available at: <https://doi.org/10.3389/fpls.2024.1459533>.
- Wazib, S., Ambardar, Y., Gautam, S., *et al.* (2025) "Asparagus racemosus (Willd.) mitigates systemic inflammation and restores angiogenic balance in L-NAME induced pre-eclampsia in experimental animals," *Journal of Ethnopharmacology*, 353, p. 120429. Available at: <https://doi.org/10.1016/j.jep.2025.120429>.

- Wazib, S., Ambardar, Y., Quasimi, H., et al. (2025) "Potential therapeutic use of Indian medicinal plants for preeclampsia management," *Journal of Ayurveda and Integrative Medicine*, 16(6), p. 101218. Available at: <https://doi.org/10.1016/j.jaim.2025.101218>.
- Wiboonpun, N., Phuwapraisirisan, P. and Tip-pyang, S. (2004) "Identification of antioxidant compound from *Asparagus racemosus*," *Phytotherapy Research*, 18(9), pp. 771–773. Available at: <https://doi.org/10.1002/ptr.1526>.
- Yadav, P. et al. (2025) "A Standardized *Asparagus Racemosus* Root Extract Improves Hormonal Balance and Menstrual Health and Reduces Vasomotor Symptoms in Perimenopausal Women: A Randomized, Double-Blind, Placebo-Controlled Study," *Journal of the American Nutrition Association*, 44(8), pp. 754–764. Available at: <https://doi.org/10.1080/27697061.2025.2510474>.
- Yuan, H. et al. (2017) "How Can Synergism of Traditional Medicines Benefit from Network Pharmacology?," *Molecules*, 22(7), p. 1135. Available at: <https://doi.org/10.3390/molecules22071135>.
- Zhang, G. et al. (2013) "Network Pharmacology: A New Approach for Chinese Herbal Medicine Research," *Evidence-Based Complementary and Alternative Medicine*, 2013, pp. 1–9. Available at: <https://doi.org/10.1155/2013/621423>.
- Zhang, H. et al. (2019) "Phytochemical compounds and biological activity in *Asparagus* roots: a review," *International Journal of Food Science & Technology*, 54(4), pp. 966–977. Available at: <https://doi.org/10.1111/ijfs.13993>.
- Zhang, W. et al. (2023) "Integration of high-throughput omics technologies in medicinal plant research: The new era of natural drug discovery," *Frontiers in Plant Science*, 14, p. 1073848. Available at: <https://doi.org/10.3389/fpls.2023.1073848>.